



I am having a Total Shoulder Replacement.

Total Shoulder Joint Replacement is a well-established surgery for restoring comfort and function to the arthritic shoulder. Shoulder arthritis is a condition in which degeneration, injury or inflammation destroys the normally smooth cartilage on the ball and socket of the shoulder joint. In a shoulder replacement surgery the damaged joint is replaced with a new artificial joint. The head of the humerus is removed and replaced with a metal prosthetic ball. The prosthesis continues down into the shaft of the humerus. The glenoid is replaced with a plastic disc.

If your shoulder has severe rotator cuff problems, a reverse shoulder replacement would be appropriate surgery. This involves replacing the humeral head with a socket and a ball as the glenoid replacement.



What happens before my surgery?

Planning ahead is the key to having less stress and achieving the best outcome from your surgery. Recovering from joint replacement surgery takes time. You can take steps before surgery that will help make your recovery easier and faster. Having a positive attitude and looking forward to getting back a good quality of life is an important part of the recovery process. The pain and deterioration of your joint has affected your daily life so just think about how much things will improve after surgery.

Before your surgery, many people will be asking about your insurance coverage, medical history, and legal arrangements. You may feel that you are answering the same questions over and over again. If you have everything written down, you can reduce your frustration and speed the process.

You should be in the best possible health before your surgery. The physical preparations you can make, can affect both the outcome of the surgery, and your recovery time. Eat well and start some light exercise. If you are overweight your doctor may suggest that you lose weight. Losing weight will help reduce stress on your new joint. If you are diabetic make sure you have clear instruction from your doctor about your medication. If you take **Blood Thinning** medication please let your surgeon know.



If you smoke, it is highly recommended that you quit or cut down, because smoking can change blood flow patterns and delay healing and slow recovery. If you drink, limit the amount of alcohol for at least 48 hours before surgery. If you are more than a social drinker please let your surgeon know to reduce the incidence of withdrawals.

Avoid any activities where you may get cuts and scratches on your arms e.g. gardening, and any of those last minute little jobs. It is very important that you have no signs of cuts, scratches or infections on your skin. If you have any sign of infection anywhere in your body, or areas of skin that are red, swollen painful or hot; cough or sore throat, contact the hospital or surgeon as this may affect your planned surgery.

The more you know about your surgery, the better you will be able to face the challenges and changes that having a Total Shoulder Replacement will make in your life. Don't ever hesitate to ask questions, voice concerns, or speak up when you do not understand.

Dr Iliesa Beci
MBChB, FRACS, FAOrthA

Dr Alan Loch
BSc, MBBS, FRACS, FAOrthA

Dr Andrew Whittle
MBBS (Hons), FRACS, FAOrthA

Dr Leo Zeller
MBBS (Hons), FRACS, FAOrthA



Planning for Extra Help:

If you live alone or have special needs, now is the time to get organised. Arrange for someone to take you to hospital and pick you up when you are ready to go home. The hospitals have a discharge by 10:00am policy on the day that you are ready to leave. You will be in a sling for after surgery, so you will need someone to look after you during the first 4-6 weeks. You will need help with transport, showering, cooking, laundry and shopping.

Planning your Food:

Prepare meals ahead of time and store in the freezer for simple reheating, or stock up on frozen dinners. Make sure you have foods that can be easily prepared and are easy on your stomach, such as soups, crackers, and soda. Move high use kitchen items so they are within easy reach. This will help reduce the need to bend and lift.

Planning your Medications:

If you take regular medication make sure your scripts are up to date. Have a written detailed list of medication, (scripted and over the counter), dosages and when you take your medication. This will enable the admission nurses to easily chart your medication, to ensure you continue taking your medication as prescribed.

Preparing your home to make it a safer place:

Your mobility and function will be restricted when you first return home. Adapt your environment before you go into hospital. Have suitable seating at home for both lounge and dining areas, at an appropriate height and with armrests. Minimise fall and trip hazards by picking up loose rugs, mats and cables. Arrange furniture so that pathways are not cluttered. Avoid slippers or shoes with open backs as they do not provide adequate support and can lead to slips and falls. Ensure toiletries, towels, toilet paper and frequently worn clothes can be reached without bending or stretching too far.

If you have stairs both inside and outside your home you will need to let your physiotherapist know so they can best guide you to manage. You may need to think about moving your bed to the ground floor of your home during this time.

If you prepare your house before going to hospital it will take the stress out of the return to home phase of your surgical journey.

Hospital Admission Process

As your admission is planned and to ensure that it is quick and easy, you must first register your details with the Hospital well in advance of your planned admission date, through the Admission Call Centre. Make sure you have all the relevant information at hand e.g. Medicare, Private Health Care provider. The Admission Call Centre will then contact you closer to the date of admission to acknowledge your admission and confirm health fund details. You will then be contacted by an admission nurse to obtain your medical history and provide you with information about your impending procedure and offers you the opportunity to ask questions about what to expect during your stay and remove the fear of the unknown.

Anaesthetic Pre-Admission Clinic

Your surgeon may require that you visit the preadmission clinic, prior to your surgery to have a health assessment. This would be carried out by an anaesthetist. They take a detailed medical history to ensure that you are in the best medical position for your anaesthetic and surgery. This may include having routine laboratory test e.g. Blood tests, urine tests, an ECG (electro-cardiogram) to monitor your heart function, and a chest X-ray may be ordered to confirm that you're fit for surgery.



What happens the day before my surgery?

The days before coming into hospital for your surgery are usually busy. Use this check list to make sure you don't forget anything.

- Bring your Orthopaedics Toowoomba surgical folder as it contains your Consent Form completed by you and your surgeon.**
- All X Rays and Scans**
- Your Medicare Card, Pension Card and Health Benefits Card.
- Your Pharmacy Benefits/Safety Net Card/Work Cover/Third party claim details/DVA Card.
- Your current medications in their pharmacy dispensed containers and a clearly written list of your medications, dosage and how often you take them. Do not bring Webster Packs.

The hospital will notify you with an admission time, where to arrive and a fasting time on this day. Fasting means no food or fluids after the time given. If you are required to take medication, take with a sip of water only.

- Shower prior to coming into hospital. Do not wear talcum powder, deodorant, perfumes or nail polish.
- Wear comfortable clothes that are easy to remove.
- Do not wear jewellery or bring valuables with you to hospital.
- Bring aids you usually require e.g. glasses, hearing aids, C Pap or walking aids in good working order.
- Do bring a hospital bag. Some items you should include are:
 - A pair of comfortable, sturdy bedroom slippers with non-skid soles.
 - A knee-length robe or gown.
 - Night attire or comfortable clothing at least 3 sets.
 - Personal care toiletries.
 - Something to read.
 - A loose fitting sweat suit or leisure wear and comfortable shoes to wear home.

If you have not asked others for help yet, do so now. Have someone check in with you daily. You'll recover more quickly if you have help instead of straining and trying to do it all yourself.

What happens during my surgery?

On the day of your surgery, arrive at the designated admission area of the relevant hospital in a timely manner. You will be greeted by the clerical staff to finalise all the information you had previously given the preadmission clinic staff. Once that has been completed you will then meet the nursing admission staff.

The nursing staff will take your blood pressure, temperature, weight, height and attend to any other tests you may require as part of your pre-surgery work up e.g. ECG, Blood Tests. They will also prepare your operative shoulder according to the surgeon's preference by painting and wrapping it in a paper towel. The nurses will also check your consent form and X-Rays. Once this has all been completed you will be changed into surgical attire and taken to a waiting room. Depending on where you are on the Surgeons operating list and how the surgery is proceeding your wait may be longer than you anticipate therefore bring a book or magazine with you.

When it is close to your operative time the theatre staff will collect you to escort you to the pre-anaesthetic bay. Here you will meet the Surgical Team.

Surgical Team

The surgical team consists of your Surgeon, his assistant and the Anaesthetist. The Surgeon will see you in the pre-anaesthetic bay before you are taken into the theatre to place an identification mark on the shoulder you have consented to have replaced. The Anaesthetist will introduce himself to you and go through your health history, medications to enable him to make a decision on the most suitable anaesthetic for you and the most appropriate pain relief regime for you. It is at this stage that the Anaesthetist will insert a drip to give you fluids while you are having your operation and to administer drugs.

During this phase you will be asked repeatedly, your name, date of birth, allergies and which side you want the operation performed. This is part of the Operating Room requirements of safe surgery. Once everything is organised you will be wheeled into the operating theatre and positioned on the operating table where a final check of your details and the operation you wish to have will occur.

Recovery

After your surgery you will be taken to the recovery room where your vital signs will be monitored closely for approximately an hour before you are taken up to the ward. At this stage your pain relief will commence according to your surgeons orders.

What happens while I am in hospital?

Day of Surgery Post-Operative Care

Once you have returned to the ward the monitoring of your vital signs will continue. You will be given the ordered pain relief that is suited for you. You will be encouraged to deep breath and cough. Your shoulder will have a dressing on it and your arm will be in a sling for immobilisation. The IV therapy will still be in your arm to enable you to receive antibiotics and pain relief; this will come out the next day if you are feeling well enough. You will be able to eat and drink if you desire. Depending on the time of day you return to the ward you may be visited by the physio department for the commencement of your therapy.

1st Day

The nurses will continue to monitor your vital signs closely during this day. If you are feeling well enough you may wish to get up for a shower. If you have a PCA for pain relief it will be removed during this morning and you will commence on a multimodal pain relief regime. If you are tolerating fluids and food the IV therapy will also be removed.



If you have not seen the physio therapist, today will be the day they get you up and start moving, doing exercises.

During this day make sure you stay on top of your pain; keep taking medication even though you may be feeling OK. As you are taking different medications for pain relief your bowel habit may change so the nurses may commence you on an aperient to keep regular. You will also commence anticoagulant therapy to prevent you getting clots.

2nd Day

You will still have your vital signs taken. Analgesia, anticoagulant therapy and bowel regime continue. The sessions with the physiotherapy department continue. The nurses will check your dressings and they may be changed if required.

3rd Day

The routine of the past two days will continue. Depending on your progress you may be ready for discharge today, if not independence will be encouraged and physiotherapy. If you require transport home today is the day to make sure that it is arranged.

4th Day

You will be discharged today, unless there is a complication from surgery. Your dressing will be changed before you go home and it is to remain in place until you see your surgeon at the postop visit 10-14 days after your surgery. The physiotherapist will see you once more before you are discharged. Discharge is usually about 10am. Make sure that you have your returning home transport organised.

What happens when I go home?

Managing Pain

After surgery, your doctor will make every effort to control your pain; however you should expect to feel some discomfort.

Many types of medicines are available to help control pain, including opioids, simple analgesics, and non-steroidal anti-inflammatory drugs (NSAIDs). Treating pain with medications can help you feel more comfortable, which will help your body heal and recover from surgery faster. So don't feel you need to 'tough it out'. When you feel less pain, you can start moving sooner and get your strength back more quickly.

Unless otherwise instructed, simple analgesia e.g. Paracetamol and NSAIDs are preferred to more powerful analgesics. Both paracetamol and NSAIDs are effective when it comes to relieving mild to moderate pain. They both relieve pain but work in different ways.

The key to successful pain relief is taking it before it is required (pre-emptive) and regularly in the first days postop. It is also ideal to take a small amount of several safe pain killers (multimodal). It is generally safe for an adult to take 2 standard Paracetamol every four hours **not exceeding 8 tablets in a day** and 2 standard NSAIDs e.g. Ibuprofen, Advil or Neurofen, every 6 hours, **not exceeding 6 tabs in a day**.



If breakthrough pain is a problem, analgesics such as Targin, Panadeine Forte, Endone and Lyrica may be prescribed.

Example of Medication Regime:

7:00a.m. – 2 NSAIDs with breakfast

9:00a.m. – 2 Panadol/ Targin/ Lyrica

1:00p.m. - 2 NSAIDs and 2 Panadol with lunch

5:00p.m. – 2 Panadol

7:00p.m. – 2 NSAIDs with dinner

9:00p.m. – 2 Panadol **or** if required stronger pain killers e.g. Panadeine Forte and Targin or Endone and Lyrica

After the acute phase of post-operative period has passed you may be comfortable on medications such as Panadol Osteo or Maxi Gesic to be taken as per instructions.

Constipation is often a side effect of taking frequent pain medication. This may be overcome by changing lifestyle e.g. increased fluid intake or increase fibre intake or by taking laxatives. A combination of the two may be needed to return to and ensure a more comfortable bowel habit.

Wound Care

The surgical dressing over your wound will usually stay in place during your hospital stay. This helps your wound to stay clean and dry. The ward nurse will change the dressing before you are discharged. This dressing is to remain intact until you have your postoperative visit to your surgeon, for the removal of the skin staples.

After you have had your staples removed your wound still needs to be looked after carefully as it heals to limit scarring, avoid discomfort and to lower the risks of infections. Another waterproof dressing will be applied and this should remain in place for a further 5 days, unless instructed otherwise by your doctor. After the dressings come off, bathe with a mild soap. Do not rub or scrub and pat dry with a clean towel.

- Do not use any lotions, salves or creams on it for at least 2 weeks from having your staples removed.
- You may notice some numbness near the wound edge. This is normal, and is due to skin nerves being cut during surgery. The numbness will decrease over time.

Driving

You will need permission from your surgeon to start driving again. This is usually around 4-6 weeks after surgery, sometimes longer. It is important to follow your surgeon's advice, for your own safety and the safety of others. Driving too early may also have implications for your insurance should you have an accident. Also, avoid travelling long distances as a passenger, by car or by plane, for at least a month after your operation, as this can cause excessive swelling and discomfort.

Physiotherapy



Commit to your exercise program. To get the most benefit from your shoulder replacement surgery, rehabilitation will be very important. Think of each exercise as a stepping-stone toward improved strength, range of motion and function.



Post-Operative Appointment.

As part of your discharge planning the ward nursing staff should give you an appointment date that will be 10 to 14 days post your surgery. Please ring your surgeon's rooms to confirm this appointment in case any changes have been made. **Phone number 07 4637 1111**

What happens if I have a problem?

Notify your Doctor for a wound review if any of the following occur:

- If there is any discharge, from the wound, that is yellow, green and offensive smelling.
- If there is heavy bleeding from the incision
- The wound edges have separated.
- Redness and hardening around the incision.
- The incision is hot to touch.
- Severe pain and tenderness.

